

# Vacuum Polarization Effects in the Worldline Variational Approach to Quantum Field Theory

R. Rosenfelder (PSI)

PANIC05, October 27, 2005

with

C. Alexandrou (Cyprus), A. Schreiber (Adelaide),  
K. Barro (ETH Zürich), M. Stingl (Münster)

1. Introduction: worldline variational approach
2. Beyond the quenched approximation
3. Results
4. Summary

## 1. Introduction

### Quantum Mechanics:

operators, states in Hilbert space

Heisenberg, Schrödinger etc.

⟷

path integrals, trajectories

Dirac, Feynman

“ WAVES ”

⟷

“ PARTICLES ”

### Field Theory:

field operators, states in Fock space

Jordan, Heisenberg, Pauli etc.

⟷

worldlines  $x_\mu(t)$

Feynman ( $\sim 1950$ )

“ FIELDS ”

⟷

“ PARTICLES ”

“ second quantization ”

↓

wins !

(see textbooks)

⟷

“ first quantization ”

↑

renaissance ... from string theory (!)

Bern & Kosower (1991)

Strassler (1992) showed how to derive the Bern-Kosower rules from the particle (worldline) representation of Quantum Field Theory

### Advantages:

- a) efficient way to calculate diagrams with many legs
- b) new approximation methods for large couplings – as in the **polaron** problem : Feynman (1955)

**best** analytical method which works for **all** coupling constants !

$$\text{thick red line} = \text{thin red line} + \text{thin red line with one loop} + \text{thin red line with two loops}$$

sums (approximately) self-energy diagrams:


$$+ \text{thin red line with one loop} + \text{thin red line with two loops} + \dots$$

## Steps in the worldline variational approach:

1. Integrate out heavy particles
2. **Schwinger representation:**

$$\frac{1}{\hat{\mathcal{O}} + i0} = -i \int_0^\infty dT e^{iT\hat{\mathcal{O}}}, \quad T : \text{proper time}$$

3. Write matrix element as (**quantum mechanical**) path integral over **particle trajectory**  $x_\mu(t)$ :



4. Integrate out light particles : only possible in **quenched approximation**  $\longrightarrow$  **retarded** (two-time) action
5. **Feynman-Jensen variational principle:**  $\langle e^{-(S-S_t)} \rangle \geq e^{-\langle (S-S_t) \rangle}$   
trial action  $S_t =$  **retarded quadratic** action
6. Solve variational equations for parameters/functions in trial action


$\implies$  result includes (approximately) all self-energy and vertex corrections !

## Applications


- Full propagator in quenched scalar model  
RR & Schreiber, Phys. Rev. **D53** (1996)
- Processes with one/two external mesons  
Alexandrou, RR & Schreiber, Nucl. Phys. **A 601** , **A 628** (1998)  
deep-inelastic inclusive scattering  
Fettes & RR, Few-Body Syst. **24** (1998)
- Improved (anisotropic) trial actions  
Schreiber & RR, Eur. Phys. J. **C 25** (2002)
- quenched QED: non-perturbative expression for anomalous mass dimension of electron  
Alexandrou, RR & Schreiber, Phys. Rev. **D 62** (2000)  
Abraham-Lorentz-like equation for electron  
RR & Schreiber, Eur. Phys. J. **C 37** (2004)
- Relativistic bound-state problem in quenched scalar model  
Barro-Bergflödt, RR & Stingl, Mod. Phys. Lett. **A 20** (2005)

Numerical solutions of var. eqs.: have found pole below  $q^2 < 4M^2$


binding energy  $\epsilon$  vs. coupling constant  $\alpha = \frac{g^2}{4\pi M^2}$  :



typical diagrams:



quenched



unquenched

## 2. Beyond the quenched approximation

Up to now **vacuum polarization** (VP) terms had to be neglected because

1. Number of worldlines for heavy particles is conserved (physics)
2. Meson/photon field cannot be integrated out with VP effects included (mathematics)

**example:** scalar Wick-Cutkosky model with  $\mathcal{L}_{\text{int}} = g \sum_i^N |\Phi_i|^2 \chi$

$\Phi_i$  : “nucleon” field with  $N$  species,  $\chi$  : “meson” field

functional integral over meson field  $\chi$  contains **determinant**

$$D[\chi] = \left[ \text{Det} \left( -\partial^2 - M_0^2 - 2g\chi \right) \right]^{-N/2}$$

**How to include VP effects?**



First possibility: expand

$$\begin{aligned} \frac{D[\chi]}{D[0]} &= \exp \left\{ -\frac{N}{2} \text{Tr} \ln \left[ 1 + \frac{2g}{\partial^2 + M_0^2} \chi \right] \right\} \\ &\approx \exp \left\{ -\frac{N}{2} \text{Tr} \left[ 2g \frac{1}{\partial^2 + M^2} \chi - \frac{(2g)^2}{2} \frac{1}{\partial^2 + M^2} \chi \frac{1}{\partial^2 + M^2} \chi + \dots \right] \right\} \end{aligned}$$

to second order in the meson field ( $M_0 \approx M$ )

**linear** terms in  $\chi$ : tadpoles (irrelevant)


**quadratic** terms in  $\chi$ :

- meson mass renormalization:  $m_0 \rightarrow m$
- modification of meson propagator by (renormalized) **one-loop VP** contribution

$\Rightarrow$   $n$ -particle **worldline** action

$$S = \underbrace{\sum_{i=1}^n \int_0^{T_i} dt \left( -\frac{1}{2} \dot{x}_i^2 \right)}_{\text{free action}} - \frac{g^2}{2} \underbrace{\sum_{i,j=1}^n \int_0^{T_i} dt \int_0^{T_j} dt'}_{\text{retarded interaction}} \int \frac{d^4 k}{(2\pi)^4} \frac{\exp[-ik \cdot (x_i(t) - x_j(t'))]}{k^2 - m^2 - \pi_r(k^2)}$$

Contains VP insertions in all meson lines:




main effect :

$$\alpha \longrightarrow \alpha^* = \frac{\alpha}{1 + \frac{\alpha}{6\pi} N}$$



↑  
number of flavors

but **no** interaction of pair-produced particles:




### 3. Results

Anomalous moment vs. coupl. constant for different  $N$  :



Binding energy vs. coupl. constant for different  $N$  :



## Second possibility:

“hybrid” approach: apply variational principle for **both** meson field  $\chi$  and worldline  $x(t)$

as in the **linear polaron model** Bogoliubov (1978)

modified meson propagator



$S_t[x, \chi] =$  quadratic in  $x(t)$  + quadratic in  $\chi$

$$+ \int_0^T dt \int d^4k \, \ell(k^2) ik \cdot x(t) \chi(k)$$



variational coupling function

**solvable** trial action, averages can be evaluated,  
(numerical) results in progress

## 4. Summary

### Worldline variational methods for field theories

- are successful due to reduction in number of variables  
e.g.  $\Phi(x), \chi(x) \longrightarrow x(t)$
- have been applied to scalar and fermionic theories  
give gauge-invariant results
- offer a new approach to the relativistic bound-state problem which includes self-energy effects and vertex corrections consistently
- can include vacuum-polarization effects

### Possible extensions:

- improved (anisotropic) trial action
- systematic second-order corrections to variational result
- inclusion of other internal degrees of freedom (color) in worldline formalism.  
QCD ?
-